



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Statistical reports of States and cities of the United States—Yearly and monthly.

IOWA—*Dubuque*.—Month of April, 1901. Census population, 36,297. Total number of deaths, 37, including enteric fever, 1; scarlet fever, 1; whooping cough, 1, and 3 from tuberculosis.

MASSACHUSETTS—*Fitchburg*.—Month of April, 1901. Census population, 31,531. Total number of deaths, 34, including smallpox, 1, and 2 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended May 25, 1901, from 91 observers, indicate that inflammation of bowels, cholera morbus, remittent fever, and scarlet fever were more prevalent and intermittent fever and smallpox less prevalent than in the preceding week. Cerebro-spinal meningitis was reported present at 4, whooping cough at 19, diphtheria at 17, enteric fever at 31, measles at 34, scarlet fever at 70, phthisis pulmonalis at 127, and smallpox at 176 places.

OHIO—*Cleveland*.—Month of April, 1901. Estimated population, 395,000. Total number of deaths, 493, including diphtheria, 7; enteric fever, 14; scarlet fever, 3; whooping cough, 1; smallpox, 6, and 41 from tuberculosis.

Columbus.—Month of April, 1901. Estimated population, 140,000. Total number of deaths, 130, including scarlet fever, 6; whooping cough, 2, and 24 from tuberculosis.

Toledo.—Month of April, 1901. Estimated population, 150,000. Total number of deaths, 149, including diphtheria, 5; enteric fever, 5; scarlet fever, 1, and 17 from tuberculosis.

WASHINGTON—*Seattle*.—Month of March, 1901. Estimated population, 90,000. Total number of deaths, 74, including diphtheria, 1; measles, 1, and 10 from tuberculosis.

Month of April, 1901. Total number of deaths, 77, including enteric fever, 2; scarlet fever, 2, and 6 from tuberculosis.

WISCONSIN—*Milwaukee*.—Month of April, 1901. Estimated population, 295,000. Total number of deaths, 323, including diphtheria, 9; enteric fever, 5; scarlet fever, 1, and 42 from tuberculosis.